```
FILE 'MEDLINE, EMBASE, BIOSIS, CAPLUS' ENTERED AT 17:55:37 ON 11 SEP 2007
           1887 S KOENIG S?/AU OR VERI M?/AU
L1
             37 S L1 AND FC
L2
L3
             17 DUP REM L2 (20 DUPLICATES REMOVED)
L4
             17 SORT L3 PY A
             0 S (CD32 OR (FC (A) GAMMA (A) RII)) (S) ANTIBOD
L5
           1503 S (CD32 OR (FC (A) GAMMA (A) RII)) (S) ANTIBOD?
L6
            250 S L6 AND INHIBITO?
L7
           117 DUP REM L7 (133 DUPLICATES REMOVED)
r_8
           2658 S 2B6 OR 3H7
L9
            618 S L9 AND ANTIBOD?
L10
              9 S L10 AND CLONE
L11
             6 DUP REM L11 (3 DUPLICATES REMOVED)
L12
L13
             35 S L9 AND CLONE
L14
             21 DUP REM L13 (14 DUPLICATES REMOVED)
```

=>

# **EAST Search History**

Ref #	Hits	Search Query	DBs	Default Operator	Plurals	Time Stamp
L1	1	"20010036459".pn.	US-PGPUB; USPAT	OR	ON	2007/09/11 17:44
L2	8039	(KOENIG).inv. OR (VERI).inv.	US-PGPUB; USPAT; USOCR; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2007/09/11 17:45
L3	54	L2 and Fc	US-PGPUB; USPAT; USOCR; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2007/09/11 17:45
L4	3348	(CD32 or (Fc adj gamma adj R)) same antibod\$	US-PGPUB; USPAT; USOCR; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2007/09/11 17:46
L5	461	(CD32 or (Fc adj gamma adj RII)) same antibod\$	USPAT	OR	ON	2007/09/11 17:47
L6	8	anti-CD32	USPAT	OR	ON	2007/09/11 17:47
L7	965	2B6 or 3H7	US-PGPUB; USPAT; USOCR; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2007/09/11 17:48
L8	81	L7 same antibod?	US-PGPUB; USPAT; USOCR; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2007/09/11 17:48



Your <b>Discoveries</b> Begin with	11_8
Begin with	US.

Search Catalog: |--- Choose Option

| Home | Ordering Info | Quick Order | Cart | Tech Support | Contact Us

Licensing **Products** Services | About ATCC |

### **Product Description**

Before submitting an order you will be asked to read and accept the terms and conditions of ATCC's Material Transfer Agreement or, in certain cases, an MTA specified by the depositing institution.

Customers in Europe, Australia, Canada, China, Hong Kong, India, Japan, Korea, Macau, Mexico, New Zealand, Singapore, and Taiwan, R.O.C. must contact a local distributor for pricing information and to place an order for ATCC cultures and products.

Cell	Biology
ATCC®	Number:

Order this item

Price:

\$203.00

**Designations:** 

Daudi

CCL-213™

Depositors:

G Klein

**Biosafety Level:** 2 [Cells Contain HERPESVIRUS]

Shipped:

frozen

Medium & Serum:

See Propagation

Growth

suspension

**Properties:** 

Organism:

Homo sapiens (human)

Morphology: lymphoblast

Source:

Organ: peripheral blood Cell type: B lymphoblast Disease: Burkitt's lymphoma

Permits/Forms: In addition to the MTA mentioned above, other ATCC and/or regulatory permits may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please <u>click here</u> for information regarding the specific requirements for

shipment to your location.

## Related Cell Culture Products

Isolation:	Isolation date: May, 1967
Applications:	transfection host (Roche FuGENE® Transfection Reagents)
Receptors:	complement, expressed Fc, expressed
Tumorigenic:	Yes, in agarose Yes, in nude mice
Reverse Transcript:	negative
DNA Profile (STR):	Amelogenin: X,Y CSF1PO: 12 D13S317: 11,12 D16S539: 10,12 D5S818: 8,13 D7S820: 8,10 THO1: 6,7 TPOX: 8,11 vWA: 15,17

Cytogenetic Analysis:	Male human karyotype with stemline number of 46. The karyotype is diploid in 66% of the cells and is stable within the stemline.
Isoenzymes:	G6PD, B
Age:	16 years
Gender:	male
Ethnicity:	Black
Comments:	The Daudi line was derived from a 16-year-old Black male with Burkitt's lymphoma by E. Klein and G. Klein in May, 1967. The cells are negative for beta-2-microglobulin. They are positive for EBNA, VCA and Surface immunoglobulin (sIg+). The line carries Epstein-Barr virus. The Daudi is a well characterized B lymphoblast cell line which has been employed extensively in studies of mechanisms of leukemogenesis.
Propagation:	ATCC complete growth medium: RPMI 1640 medium with 2 mM L-glutamine adjusted to contain 1.5 g/L sodium bicarbonate, 4.5 g/L glucose, 10 mM HEPES, and 1.0 mM sodium pyruvate, 90%; fetal bovine serum, 10% Temperature: 37.0C Atmosphere: air, 95%; carbon dioxide (CO2), 5%
Subculturing:	Protocol: Cultures can be maintained by the addition of fresh medium or replacement of medium. Alternatively, cultures can be established by centrifugation with subsequent resuspension at 3 to 5 X 10(5) viable cells/ml.  Interval: Maintain cell density between 3 X 10(5) and 2 to 3 X 10(6) viable cells/ml.  Medium renewal: Add fresh medium every 2 to 3 days (depending on cell density)
Preservation:	Freeze medium: Complete growth medium supplemented with 5% (v/v) DMSO Storage temperature: liquid nitrogen vapor phase
Related Products:	Recommended medium (without the additional supplements or serum described under ATCC Medium): ATCC 30-2001 recommended serum: ATCC 30-2020
References:	22550: Ohsugi Y , et al. Tumorigenicity of human malignant lymphoblasts: comparative study with unmanipulated nude mice, antilymphocyte serum-treated nude mice, and X- irradiated nude mice. J. Natl. Cancer Inst. 65: 715-718, 1980. PubMed: 6932523 23017: Klein E , et al. Surface IgM-kappa specificity on a Burkitt lymphoma cell in vivo and in derived culture lines. Cancer Res. 28: 1300-1310, 1968. PubMed: 4174339 26046: Huber C , et al. Surface receptors on human haematopoietic cell lines. Clin. Exp. Immunol. 25: 367-378, 1976. PubMed: 963908 26047: Nilsson K , et al. Tumorigenicity of human hematopoietic cell lines in athymic nude mice. Int. J. Cancer 19: 337-344, 1977. PubMed: 14896 28315: Gao Y , et al. Induction of an exceptionally high-level, nontranslated, Epstein-Barr virus-encoded polyadenylated transcript in the uurkitts lymphoma line Daudi. J. Virol. 71: 84-94, 1997. PubMed: 8985326 32286: Cuthbert JA , Lipsky PE . Regulation of proliferation and Ras localization in transformed cells by products of mevalonate metabolism. Cancer Res. 57: 3498-3504, 1997. PubMed: 9270019 32830: Yamaguchi Y , et al. Biochemical characterization and intracellular localization of the Menkes disease protein. Proc. Natl. Acad. Sci. USA 93: 14030-14035, 1996. PubMed: 8943055 33091: Lewis JA , et al. Inhibition of mitochondrial function by interferon. J. Biol. Chem. 271: 13184-13190, 1996. PubMed: 8662694 33115: Montoya JG , et al. Human CD4+ and CD8+ T lymphocytes are both cytotoxic to Toxoplasma gondii-infected cells. Infect. Immun. 64: 176-181, 1996. PubMed: 8557337

#### **Notices and Disclaimers**

ATCC products are intended for laboratory research purposes only, unless noted otherwise. They are not intended for use in humans.  $\dot{}$ 

While ATCC uses reasonable efforts to include accurate and up-to-date information on this site, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

All prices are listed in U.S. dollars and are subject to change without notice. A discount off the current list price will be applied to most cultures for nonprofit institutions in the United States. Cultures that are ordered as test tubes or flasks will carry an additional laboratory fee. Fees for permits, shipping, and handling may apply.

You may continue your ATCC Number search by typing in your search criteria below or returning to the <u>ATCC Search Page</u>. For more information please review the <u>Search Help</u>.

CCL-213

ATEC Number Search

Clear Search

#### **Home Page Archive**

Home Ordering Info Quick Order Support About ATCC Contact Us
Privacy Policy Terms of Use ATCC MTA

© 2007 American Type Culture Collection (ATCC).
All rights reserved.